

Maternal Vaccination



Resources for healthcare professionals

Vaccines help keep your pregnant patients and their growing families healthy.

Last Updated September, 2016

Vaccine	Before pregnancy	During pregnancy	After pregnancy	Type of vaccine
Influenza	Yes	Yes, during flu season	Yes	Inactivated
Tdap	May be recommended; it is better to vaccinate during pregnancy when possible	Yes, during each pregnancy	Yes, immediately postpartum, if Tdap never received in lifetime; it is better to vaccinate during pregnancy	Toxoid/ Inactivated
Td	May be recommended	May be recommended, but Tdap is preferred	May be recommended	Toxoid
Hepatitis A	May be recommended	May be recommended	May be recommended	Inactivated
Hepatitis B	May be recommended	May be recommended	May be recommended	Inactivated
Meningococcal	May be recommended	Base decision on risk vs. benefit; inadequate data for specific recommendation	May be recommended	Inactivated
Pneumococcal	May be recommended	Base decision on risk vs. benefit; inadequate data for specific recommendation	May be recommended	Inactivated
HPV	May be recommended (through 26 years of age)	No	May be recommended (through 26 years of age)	Inactivated
MMR	May be recommended; once received, avoid conception for 4 weeks	No	May be recommended	Live
Varicella	May be recommended; once received, avoid conception for 4 weeks	No	May be recommended	Live

For more information, visit: www.cdc.gov/vaccines/pregnancy

Get an answer to your specific question by e-mailing cdcinfo@cdc.gov or calling 800-CDC-INFO (232-4636)



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention